

**DESCRIPTION OF THE MINT LEAF BEETLE, *CHRYSOLINA HERBACEA*  
(DUFTSCHMID, 1825) (COLEOPTERA: CHRYSOMELIDAE) FROM KURDISTAN  
REGION– IRAQ**

**Hani A. Ibrahim<sup>1</sup>, Zewar Z. Omar<sup>2\*</sup>, Gazang T. Omar<sup>3</sup> and Nabeel A. Mawlood<sup>3</sup>**

<sup>1</sup>Department of Horticulture, Technical College, Akre/ Duhok Polytechnic University, Duhok,  
Iraq

<sup>2</sup> Department of Plant Protection, Khabat Technical institute, Erbil Polytechnic University, Erbil,  
Iraq

<sup>3</sup>Department of Plant Protection, College of Agricultural Engineering Science, Salahaddin  
University, Iraq

\*Corresponding author email: [zewar.omar@epu.edu.iq](mailto:zewar.omar@epu.edu.iq)

**ABSTRACT**

The present work includes a detailed description of the mint leaf beetle, *Chrysolina herbacea* (Duftschmid) from Kurdistan Region– Iraq. The specimens were collected from the mint leaf during January to June of 2021. The taxon is easily distinguishable, where the labrum is dark green, and transverse oval shaped, slightly emarginated at the middle. Mandibles are asymmetrical, and apical part of left mandible is bidenticates. Fourth segment of maxillary palps cylindrical, as long as third segment. Antenna dark brown and filiform, reaching to the 1/3 of elytra base. Tegmen distinctly Y-shaped. Median lob is sword shaped and acute at the apical part.

Keywords: Coleoptera, Description, *Chrysolina herbacea*, Chrysomelidae, Kurdistan Region, Iraq

**INTRODUCTION**

Chrysomelidae commonly is known as leaf beetles and are the second largest family of phytophagous Coleoptera, and it is widely distributed worldwide with over 55,000 species in 17 families and more than 2,500 genera (Cox,1976; Booth *et al.*, 1990; Gruev, 1992). Adult and larval feed on all sorts of plant tissue, and all species are fully herbivorous. Many are serious pests to cultivated plants, and some of them are regarded as pests in agriculture and forestry (Warchałowski 2003; Özdikmen and Aslan, 2009). *Chrysolina* Motschulsky, 1860 is a very large and diverse genus of the family. Most species are distributed in Europe, Asia, with a small number of species inhabiting North America and were introduced to Australia and Africa (Bienkowski, 2001). Kippenberg (2001) reported that the genus consists of approximately 345 species in palearctic region. Bechyné (1952) explained a taxonomical review of the genus and described some new subgenera and species. The genus *Chrysolina* is a very diverse of leaf-beetles consist of nearly 450 species belonging to 65 subgenera that have been identified (Bieńkowski, 2001), and new species are still being described (Ge *et al.*, 2011; Bourdonné *et al.*, 2013; Lopatin, 2014). *Chrysolina herbacea* (Duftschmid) is one of very important species of the genus, as it is a monophagous insect feeding on various Lamiaceae (Mint, pennyroyal), with a preference for the mint plants (Çağrı *et al.*, 2015). Both larvae and adult of the beetles feed on mint

leaves (King, 2015). In Iraq, El-Haidari *et al.*, (1972) recorded three species of the family including *Chrysolina marginata*. Al-Ali (1977) reported 17 species of the family involving two species that belong to the genus; which are *Chrysolina chalcites* (Germ.) and *Chrysolina grata* (Falderman).

The aim of this paper is a detailed description of mint leaf beetle, *Chrysolina herbacea* (Duftschmid) which collected from mint leaf in different areas of Kurdistan Region– Iraq.

## MATERIALS AND METHODS

The present paper is based on 50 specimens. The samples of leaf beetle, *Chrysolina herbacea* (Duftschmid) were collected during January to June in some areas of Erbil and Sulimani governorates including (Qushtapa and Benslaw) and (Kalar and Beckrajo); respectively, in Kurdistan region - Iraq.

The specimens were placed in boiling water for 10-15 minutes to soften their parts. Then the parts were separated under dissecting microscope to three parts; head, thorax and abdomen. The head and abdomen were soaked in a beaker contained 10% KOH and placed on fire with shaking for about (10-15) minutes for dissolving of lipids materials of the body and destroying the muscles. After that, the samples were placed in distilled water for 2-3 minutes in order to neutralize the alkali. The parts were placed in ethyl alcohol 25% and dissected under microscope then transferred to ethyl alcohol 50%, 75% and 100%, respectively for two minutes of each concentration for water dehydration. The samples were placed in xylol for two minutes, for translucency. Finally, the parts finely were placed on slides with a drop of DPX solution and covered by cover slides to be prepared slides for examination (Lane and Grosskey, 1993; Mawlood *et al.*, 2016; Abdulla *et al.*, 2020).

The specimens were deposited in the Insects Museum in the Department of Plant protection- College of Engineering Sciences- University Salahaddin-Erbil Governorate Kurdistan region- Iraq.

## RESULTS and DISSCUSION

### *Chrysolina herbacea* (Duftschmid, 1825)

Synonyms

*Chrysomela herbacea* (Duftschmidt, 1825)

*Chrysomela menthastri* (Suffrian, 1851)

*Chrysomela fulminans* (Suffrian, 1851)

## DESCRIPTION

### Body

Oval, strongly convex and shiny metallic green, bronze color with black legs and antennae. Length 6.1-7.5 mm, width 3.7-4.4 mm.

### Head

Spherical shaped, metallic green with narrow sparsely setose and by punctate. Eyes black oval strongly prominent. Frons shine green, slightly convex, with moderately dense of punctures near clypeus. Clypeus triangular, shiny black green sparsely punctate. Fronto-clypeal suture smooth. Gena brown, sparsely punctate. Labrum (Fig.1A) dark green, posterior margin slightly emarginated at the middle, surface sparsely setose and punctate. Mandibles asymmetrical, high sclerotized, left mandible (Fig.1B) triangular, apical with two denticles, inner denticles shorter

than the outer, right mandible (Fig.1C) resembling to left mandible except, the apical part with single denticles. Maxilla (Fig.1D) dark brown, high sclerotized; basigalea elongated oval slightly longer than distigalea, apical part of each with dense of long yellow setae, 1<sup>st</sup> -3<sup>rd</sup> segments of maxillary palps cup shaped, sparsely dark brown setose, 2<sup>nd</sup> segment 1.2 as long as 3<sup>rd</sup> segment, 4<sup>th</sup> segment broadly oval, obliquely truncate, as long as third segment. Labium dark brown, 2<sup>nd</sup> segment of labial palp cup shaped, 1.2 times as long as 3<sup>rd</sup> segment, which are oval. Antenna (Fig.1E) dark brown consist of 11 antennomeres, 2<sup>nd</sup> antennomere elongated oval, 2.1 times as long as the pedicel and subequal to the 3<sup>rd</sup> antennomere, 3<sup>th</sup> -5<sup>th</sup> antennomeres nearly tubular shaped, 3<sup>th</sup> antennomere 1.1 times as long as the 4<sup>th</sup> once, 6<sup>th</sup> -10<sup>th</sup> segments cup shaped, 10<sup>th</sup> segment 1.1 times as long as 9<sup>th</sup> segment, 11<sup>th</sup> segment elongated oval 1.2 times as long as 9<sup>th</sup>.

### Thorax

Prothorax shiny metallic green, nearly 1.75x longer than wide and narrower than the elytral base. pronotum convex, shiny black green, transverse, nearly twice longer than wide, anterior and posterior margins moderately curved. Pronotal disc sparsely and unevenly punctured with short pale yellow setae. Lateral margins of pronotum slightly curved, anterior and, posterior angles moderately produced, basal angles sharp. Scutellum triangular, anterior margin straight, apical part rounded, lateral margin curved, surface with high dense of granules. Anterior margin of prosternum slightly straight bear row of short brown setae, prosternal process nearly globular shaped, surface with moderate dense of fine punctures. Elytra (Fig.1F) oval, convex, metallic green, length 4.0-4.6 mm, covered, sparsely punctate; internal border of the lateral margin fringed with short, fine setae near the apex. Epipleura slightly straight, surface with low dense of punctures. Hind wing (Fig.1G) red color, veins yellow, costa short, less than half of subcostal, radius veins incomplete, radial cell triangular shaped, median spur extending to wing margin, cross veins r-m and cu-an absent, veins cu1 present but undivided; vein 2A slightly reduced. Fore legs (Fig.1H) metallic green, forecoxa conical shaped; femur cylindrical with low dense of punctures and short black setae; tibia tubular shaped, slightly longer than femur with short black setae, apical part without spurse; 1/6 of apical part with row of short brown erect spines; Protarsus black consist of five segments, 1<sup>st</sup> -2<sup>nd</sup> segments cup shaped, 2<sup>nd</sup> slightly smaller than the 1<sup>st</sup> segment, 3<sup>rd</sup> segment bilobed, 4<sup>th</sup> segment very small hidden beneath third segment, 5<sup>th</sup> segment is the longest, tubular shaped, 1.9 times as long as 2<sup>nd</sup> segment, outer surface of the segments 1<sup>st</sup> - 3<sup>rd</sup> covered with high dense of very short setae, claw hook shaped, brown color. Middle legs resemble to fore legs except the coxa nearly oval. Hind legs resemble to the forelegs, except the coxa nearly plate shaped.

### Abdomen

Abdomen oval shaped, metallic green, consist of five visible sternites. The abdomen not grooved for reception of hind legs. 1<sup>st</sup> - 4<sup>th</sup> sternites rectangular, 1<sup>st</sup> sternite 1.2 times as long as the 2<sup>nd</sup> sternite, 5<sup>th</sup> sternite oval, 1.3 times as long as the 4<sup>th</sup> sternite, each sternites bear moderately dense of short pale yellow setae, with moderate dense of punctures. Tergites seven segmented, 1<sup>st</sup> - 6<sup>th</sup> tergites rectangular, anterior and posterior margin of 1<sup>st</sup> - 4<sup>th</sup> tergites are straight, posterior margin

of slightly emarginated at the middle and 2.6 times as long as 6<sup>th</sup> . pygidium(Fig.1I) nearly cup shaped low sclerotized at the middle.

### Male genitalia

Aedeagus (Fig.1K) height sclerotized, dark brown, flattened in cross-section, strongly curved dorso-ventrally. Length 1.4- 2.0 mm. Phallobase dark yellow, oval. Tegmen (Fig. 1J) height sclerotized, inverted V-shaped and partially surrounds the aedeagus. Medial lob cylindrical shaped, moderately curved, its apex is lanceolate. Endophallus long and moderately sclerotized, joined to the parameres by simple temones. Ejaculatory duct long, tubular shaped. flagellum Invisible.



Fig.1 *Chrysolina herbacea* (Duftschmid)

A. Labrum B. Left mandible C. Right mandible D. Maxilla E. Antenna F. Elytra  
G. Hind wing H. Fore leg I. Pygidium J. Tegmen K. Aedeagus (Lateral view)

Scale bar: a, b ,c ,d ,e, i, j and k = 0.5mm ; f, g, and h = 2mm

## ACKNOWLEDGMENTS

We Sincerely thank the specialist, Assist. Prof. Dr. Hanna Hani Al-safar in Iraq Natural History Research Center and Museum -University of Baghdad for their assistance to identify the species.

## REFERENCES

- Abdulla, B.S., Ahmmed, S. H., Mawlood, N. A. and Omar, Z. Z. 2020. A New record of European chafer beetles, *Amphimallon majale* (Razoumowsky, 1789) (Coleoptera: Melolonthidae) from Iraq. Plant Archives. 20 (2): 6357-6361.
- Al-Ali, A.S. 1977. Phytophagous and entomophagous insects and mites of Iraq. Natural History Research Center, Publication, 33: 142.
- Bechyné, J. 1952. Achter Beitrag zur Kenntnis der Gattung *Chrysolina* Motsch. (Col., Phytoph., Chrysomelidae). Ent. Arbeit. Mus. G. Frey, 3: 351-385.
- Bienkowski, A. J. O. 2001. A study on the genus *Chrysolina* Motschulsky, 1860, with a checklist of all the described subgenera, species, subspecies, and synonyms (Coleoptera: Chrysomelidae: Chrysomelinae) Genus - International Journal of Invertebrate Taxonomy. 12,105–235.
- Booth, R. G., Cox, M. L. and Madage, R. B. 1990. The Guides to Insects of Importance to Man, 3. Coleoptera. Printed in the UK at the University Press, 384 p.
- Bourdonné, J. C., Doguet, S. and Petitpierre, E. 2013. *Chrysolina (Stichoptera) oceanoripensis* nova species, endémique des dunes françaises des Landes de Gascogne et considérations sur le sous-genre Stichoptera Motschulsky, 1860 (Coleoptera, Chrysomelidae). Nouvelle Revue d'Entomologie, 29(1): 33–52.
- Çağrı, B., Mustafa, C., Tuba, B. and Onur, T. 2015. Role of Different Thymus Essential Oils on Feeding Performance of *Leptinotarsa decemlineata* Say, 1824 (Chrysomelidae: Coleoptera). Advances in Zoology and Botany. 3(1): 6-9.
- Cox, M. L. 1976. The taxonomy and biology of the British Chrysomelidae. PH.D. Thesis, Faculty of Agriculture, Newcastle University. 140pp.
- El-Haidari, H., Fattah, Y. M., and Sultan, J. A. (1972). Contribution to the insect fauna of Iraq. Part4. Direct. Gen. Plant prot. Iraq, Bull., 18: 17.

Ge, S., Daccordi, M., Li, W. and Yang, X. K. 2011. Seven new species of the genus *Chrysolina* Motschulsky from China (Coleoptera: Chrysomelidae: Chrysomelinae). *Zootaxa*, 2736: 31–43.

Gruev, B. A. 1992. Geographical distribution of Lamprosomatinae, Eumolpinae, Chrysomelinae, Alticinae, Hispinae and Cassidinae (Coleoptera, Chrysomelidae) on the Balkan Peninsula. Plovdiv University Press, 512p.

King, J. E. 2015. Beetle biodiversity response to vegetation restoration of mid-valley riparian woodland in the Lower Rio Grande Valley of southern Texas. PhD [dissertation], Texas A&M University College Station viii.

Kippenberg, H. 2001. Neuordnung der Gonioctenavariabilis- Gruppe (Coleoptera, Chrysomelidae). *Entomologische Blätter für Biologie und Systematik der Käfer*, 97:13-34.

Lane, R. P. and Crosskey R.W. 1993. *Medical Insects and Arachnids. The Natural History Museum.* Chapman and Hall, London.

Lopatin, I.K. 2014. New species of leaf-beetles (Coleoptera, Chrysomelidae) from China: XI. *Entomological Review*. 94(3): 397–407.

Mawlood, N.A., M.I. Hamad and Y.M. Abdullah 2016. A new record of glaphyrid scarab beetles, *Eulasia vitatta* (Fabricius, 1775) (Coleoptera, Glaphyridae) from Erbil Kurdistan region-Iraq. *Zanco Journal of Pure and Applied Sciences*, 28(3): 1-4.

Özdikmen, H. and Aslan, K. 2009. First records of some leaf beetles for Mediterranean region in Turkey and South Turkey (Coleoptera: Chrysomelidae). *Munis Entomology & Zoology*, 4 (1): 276-279.

Warchałowski, A. 2003. *Chrysomelidae: The leaf beetles of Europe and the Mediterranean area.* Natura optima dux Foundation, Warszawa, 600p.